

CONTENTS

*Special Issue: Theoretical Chemistry 1990. Proceedings of the XIXth International Congress
of Theoretical Chemists of Latin Expression, Rome, Italy, 10–14 September 1990*

<i>Editorial</i>	x
Results of the correlation energy functionals versus the wavefunction type	
E. San-Fabián, F. Moscardó, J.M. Pérez-Jordá (Alicante, Spain), A. Aguado and M. Paniagua (Madrid, Spain)	1
Modified relativistic effective potentials for transition elements	
L. Fernandez Pachos (Madrid, Spain)	13
A new procedure to obtain GTO wave functions and to analyze their quality	
J.M. García de la Vega and B. Miguel (Madrid, Spain)	21
A practical approach to the computation of the electrostatic energy in large molecules	
M. Orozco and F.J. Luque (Barcelona, Spain)	31
Bonding in the CuCH₄⁺ and the FeCO⁺ cations	
G. Berthier (Paris, France), R. Cimiraglia (Pisa, Italy), A. Daoudi (Fes-Atlas, Morocco), H. Mestdagh, C. Rolando (Paris, France) and M. Suard (Montrouge, France)	43
On the use of diffuse functions for describing dissociative electron transfer reactions	
V. Pérez, J.M. Lluch and J. Bertrán (Barcelona, Spain)	51
Recursion relations for matrix elements and Franck–Condon factors of rotating Morse oscillators	
A. López Piñeiro, M.L. Sánchez and B. Moreno (Badajoz, Spain)	63
Memory superoperator representation of projected one-particle casual propagators	
R.C. Bochicchio and H. Grinberg (Buenos Aires, Argentina)	71
Random phase approximation response of molecular systems to electromagnetic fields: a very preliminary investigation of choice criteria for Slater-type basis sets	
C. Guidotti (Pisa, Italy), U.T. Lamanna (Bari, Italy), G.P. Arrighini and N. Durante (Pisa, Italy)	77
The adiabatic multi-step separation method for vibrational states: application to SO₂	
J. Zúñiga, J.M. Francés, J.V. Lorente and A. Requena (Murcia, Spain)	91
A parameter-free adiabatic potential as a probe for electron–molecule collisions	
F.A. Gianturco (Rome, Italy)	99
The diabatic approximation in the classical frame	
P. Villarreal, S. Miret-Artés, O. Roncero and G. Delgado-Barrio (Madrid, Spain)	117
Regional virial in diatomic molecules: two criteria for defining zero-flux dividing surfaces	
V. Mujica (Caracas, Venezuela), M. García-Sucre (Caracas, Venezuela and São Carlos, Brazil) and Y. Aray (Maracaibo, Venezuela)	125
A numerical procedure to obtain accurate potential energy curves for diatomic molecules	
A. Aguado, J.J. Camacho and M. Paniagua (Madrid, Spain)	135
Improving the one-electron states of ab initio LCAO-GTO calculations in momentum space. Application to Be and B⁺ atoms	
L. De Windt, J.G. Fripiat, J. Delhalle (Namur, Belgium) and M. Defranceschi (Gif-sur-Yvette, France)	145
Relativistic effects on the computation of oscillator strengths for the principal series in Na-like systems	
C. Lavín, I. Martín (Valladolid, Spain) and J. Karwowski (Torún, Poland)	161
Electronic structure of tetrahedral tin acetylides by pseudopotential ab initio calculation and gas-phase UV photoelectron spectroscopy	
M.V. Andreucci, M. Bossa, C. Cauletti, S. Stranges (Rome, Italy), K. Horchler and B. Wrackmeyer (Bayreuth, FRG)	171
Polarization functions for the calculations of polarizability and hyperpolarizability. Application to the helium atom and the hydrogen molecule	
N. El Bakali Kassimi, M. Tadjeddine, J.P. Flament, G. Berthier and H.P. Gervais (Palaiseau, France)	177

Finite-field fourth-order MBPT calculations of dipole moments, field, and field-gradient polarizabilities of diatomic molecules G.H.F. Diercksen and A.J. Hernández (Garching b. München, FRG).....	191
Dipole moments and polarizabilities of some substituted pyridine-1-oxides for optoelectronics G. Berthier (Paris, France), M. Defranceschi (Gif-sur-Yvette, France), P. Lazzaretti (Modena, Italy), G. Tsoucaris (Châtenay-Malabry, France) and R. Zanasi (Modena, Italy).....	205
Ab initio BSSE-EICP calculations of thermodynamic properties on linear hydrogen fluoride dimerization S. Tolosa, F. Manzano, E.A. Ojalvo and F.J. Olivares del Valle (Badajoz, Spain).....	219
MCSCF determination of the KO molecule ground state. CAS and basis set dependence L. Serrano-Andrés, A. Sánchez de Merás, R. Pou-Amérigo and I. Nebot-Gil (Valencia, Spain)	229
Theoretical investigation of the electronic properties of poly(thienylene vinylenes) and poly(furylene vinylenes) E. Ortí, R. Crespo and M.C. Piqueras (Burjassot, Spain)	243
Theoretical comparison of proximate peri and ortho substituent effects on proton chemical shifts T. Herr, M.B. Ferraro and R.H. Contreras (Buenos Aires, Argentina)	253
Calculation of substituent electric fields T. Herr, M.B. Ferraro and R.H. Contreras (Buenos Aires, Argentina)	261
Ab initio study of the nitrogen lone-pair proximity effect on a methyl C–H bond in 2-methylpyridine M.F. Tufró, R.H. Contreras (Buenos Aires, Argentina) and J.C. Facelli (Salt Lake City, UT, USA)	271
Ab initio study of the electronic, vibrational and rotational properties of perfluoroacetylchloride and trifluoroacetaldehyde E. Ottavianelli, E.A. Castro and A.H. Jubert (La Plata, Argentina)	279
Conformational properties of ethanediol in aqueous solution as described by the continuous model of the solvent G. Alagona and C. Ghio (Pisa, Italy)	287
A theoretical study of the spectrum of 3d ⁿ ions adsorbed on ionic crystals. II. Co ²⁺ on calcite and aragonite A. Julg (Marseille, France) and M. Fernández-Núñez (Madrid, Spain).....	301
Semi-empirical model potentials for the interaction between two O ₂ ground state molecules B. Bussery and M. Aubert-Frecon (Villeurbanne, France)	307
Comparative ab initio and semi-empirical study of hydrogen bonded complexes of NH ₃ and H ₂ O E.L. Coitiño, O.N. Ventura and R.M. Sosa (Montevideo, Uruguay)	315
An algorithm based in Ewald's method to calculate lattice sums in the framework of crystal field theory. I. Application to the calculation of electrostatic potential inside MX ionic crystals J. Andrés, A. Beltrán, J. Monterde (Castelló, Spain) and M. Fernández-Núñez (Madrid, Spain)	329
Diatomeric energy partition as a tool for bond analysis M. Sánchez and F. Ruette (Caracas, Venezuela)	335
Molecular graphics investigation of the addition of nucleophiles to (η^4 -butadiene)M(CO) ₃ complexes (M=Fe, CO ⁺) J. Weber, P.-Y. Morgantini (Geneva, Switzerland) and O. Eisenstein (Orsay, France)	343
Influence du solvant sur la stabilité des hydroxypyrones. I H. Meghezzi et A. Boucekkine (Alger, Algeria)	359
Molecular surface calculations on organic compounds. Molecular area–aqueous solubility relationships E. Silla, I. Tuñón, F. Villar (Valencia, Spain) and J.L. Pascual-Ahuir (New York, NY, USA)	369

Dissociative adsorption process of acetaldehyde on Pt (<i>h,k,l</i>) basal surfaces. An EHMO study J.M. Pérez (Alicante, Spain), F. Cases (Alcoi, Spain), F.J. Puig and J.L. Vázquez (Alicante, Spain)	379
Theoretical modelling of molybdenum interaction with a graphite furnace surface A. Sierraalta, Z. de Benzo, W. Araujo and F. Ruette (Caracas, Venezuela)	387
Theoretical calculations for catalytic activation of N ₂ on a modelled iron adsorption site L. Rincón, F. Ruette and A. Hernández (Caracas, Venezuela)	395
Vibrational modelling in large polycyclic aromatic hydrocarbons N. Rougeau, J.P. Flament, P. Youkhribache, H.P. Gervais and G. Berthier (Palaiseau, France)	405
Molecular aggregation of polycyclic aromatic hydrocarbons. A theoretical modelling of coronene aggregation J. Rodríguez (Caracas, Venezuela), J. Sánchez-Marín, F. Torrens (Valencia, Spain) and F. Ruette (Caracas, Venezuela)	429
Theoretical study of the formation of cyclopropenyl cation from CH ⁺ and acetylene R. López, J.A. Sordo and T.L. Sordo (Oviedo, Spain)	443
Ab initio characterization of possible dissociation pathways for multiphoton ionization of the water dimer in supersonic free jets R.M. Sosa, K. Irving and O.N. Ventura (Montevideo, Uruguay)	453
Amidine decomposition mechanism. A theoretical study J. Andrés, A. Beltran, M. Carda, J. Krechl, J. Monterde and E. Silla (Castelló, Spain)	465
Ab initio calculation of protonation and lithiation energy of acetamidoxime M. Cotta Ramusino (Rome, Italy), G. La Manna (Palermo, Italy) and O. Tomagnini (Rome, Italy)	473
A theoretical study of the acidity of nitrogen heterocycles J.C. del Valle and J.L.G. de Paz (Madrid, Spain)	481
AM1 study of the cycloaddition of singlet methylene to butadiene and the vinylcyclopropane rearrangement J.J. Quirante, F. Enríquez (Málaga, Spain) and J.M. Hernando (Valladolid, Spain)	493
A proposed topography for the histamine H ₂ -receptor from structural AM1 calculations F.M.L.G. Stamato, E. Longo, M.A. Perez, M.A.P. Guimaraes, Jr. (San Carlos, Brazil) and Y.G. Smeyers (Madrid, Spain)	505
Quantum molecular similarity measures and the <i>n</i> -dimensional representation of a molecular set: phenyldimethylthiazines ¹ R. Carbó and B. Calabuig (Girona, Spain)	517
PQQ or TOPA? A theoretical suggestion for the organic cofactor of amine oxidases M. Bossa, I. Gelsomino, G.O. Morpurgo, L. Morpurgo and A. Talamo (Rome, Italy)	533
<i>Author index</i>	539
<i>Subject index</i>	541

